

INDIA AIDS INDICATOR TABLES

INDIA

Source NFHS: National Family Health Survey

Condom Availability and Quality

#	Indicator	2005-06 NFHS			
		Urban		Rural	
		M	F	M	F
2.5	Knowledge of a formal source of condoms among young people. Ages 15-24.	91.3	38.8	81	17.7

Stigma and Discrimination

#	Indicator	2005-06 NFHS			
		Urban		Rural	
		M	F	M	F
3.1	Accepting attitudes towards those living with HIV – Composite of 4 components. Ages 15-49.	41.6	36.6	33.4	31.2
3.1.1	Accepting attitudes - Willing to care for family member sick with AIDS (1) Ages 15-49.	80.9	78.1	74.6	71.9
3.1.2	Accepting attitudes - Would buy fresh vegetables from a shopkeeper with AIDS (2) Ages 15-49.	72.3	67	56.1	54.6
3.1.3	Accepting attitudes - Female teacher who is HIV+ but not sick should be allowed to continue teaching in school (3) Ages 15-49.	78.5	79.6	65.8	69.3
3.1.4	Accepting attitudes - Not secretive about family member's HIV status (4) Ages 15-49.	63.1	61	66.6	66.2

Knowledge

#	Indicator	2005-06 NFHS			
		Urban		Rural	
		M	F	M	F
4.0.1	Heard of HIV/AIDS. Ages 15-49.	94.8	83.2	77.1	50
4.1	Knowledge of HIV prevention methods-composite of 2 components (prompted) Ages 15-49.	78.5	50.5	57.1	23.5
4.1.2	Knowledge of HIV prevention methods - Use of condoms (prompted) Ages 15-49.	83.6	55.8	62.2	26.7
4.1.3	Knowledge of HIV prevention methods - Only one partner (prompted) Ages 15-49.	85.8	64.7	65.4	34.7
4.1.4	Knowledge of HIV prevention methods - Abstain from sex (prompted) Ages 15-49.	79	58.8	60.2	31.1
4.2	No incorrect beliefs about AIDS - Composite of 3 components Ages 15-49.	51.9	39.2	28.9	15.7
4.2.1	No incorrect beliefs about AIDS - Healthy-looking person can have the AIDS virus Ages 15-49.	75	56.2	53.3	28.5
4.2.2	No incorrect beliefs about AIDS - AIDS cannot be transmitted by mosquito bites Ages 15-49.	67.3	57.7	44.7	23.8
4.2.6	No incorrect beliefs about AIDS - Cannot become infected by sharing food with someone who has AIDS Ages 15-49.	77.7	63.3	51.7	31.5

INDIA AIDS INDICATOR TABLES

4.5	Knowledge of prevention of mother to child transmission of HIV Ages 15-49.	74.6	67	56.8	36.7
4.5.1	Knowledge of prevention of MTCT - Can be prevented by mother taking ART during pregnancy Ages 15-49.	24.6	28.4	17.9	13.8
4.6	Comprehensive correct knowledge about AIDS (2 ways to prevent AIDS and reject 3 misconceptions) Ages 15-49.	46.6	30.3	25.1	11
4.7	Comprehensive correct knowledge about AIDS among young people age 15-24 (2 ways to prevent AIDS and reject 3 misconceptions)	47.1	32.9	29.4	14

Voluntary Counseling and Testing

		2005-06 NFHS			
		Urban		Rural	
#	Indicator	M	F	M	F
5.1.1	Population receiving an HIV test. Ages 15-49.	5.7	5.7	2.6	2
5.6	HIV testing behavior among young people, sexually active in the last 12 months	1.9	5.3	1	1.7

Sexual Negotiation and Attitudes

		2005-06 NFHS			
		Urban		Rural	
#	Indicator	M	F	M	F
7.1	Women's ability to negotiate safer sex with husband. Ages 15-49.	93.3	80.9	86.1	76.4

Note: Percentage of women who believe that, if a husband has a STD, his wife is justified in refusing to have sexual intercourse with him

Percentage of men who believe that, if a husband has a STD, his wife is justified in refusing to have sexual intercourse with him or asking that they use a condom

Sexual Behavior

		2005-06 NFHS			
		Urban		Rural	
#	Indicator	M	F	M	F
8.1	Higher risk sex in the last year. Ages 15-49.	5.5	0.2	5.2	0.2
8.1.1	Multiple partners in the last year Ages 15-49.	1.8	0.1	2.1	0.1
8.2	Condom use at last higher risk sex. Ages 15-49.	52.9	24.9	29.7	11.7
8.2a	Condom use at last higher risk sex. Ages 15-	M: 31.3		F: 20	
8.2b	Condom use at last higher risk sex. Ages 20-	M: 40.7		F: 25.1	
8.2c	Condom use at last higher risk sex. Ages 25-29.	M: 46.4		F: (25.4)	
8.3	Commercial sex in last year. Ages 15-49.	0.8		0.6	
8.4	Condom use at last commercial sex, reported by client. Male only. Ages 15-49.	72.7		54.4	

INDIA AIDS INDICATOR TABLES

Young People's Sexual Behavior

#	Indicator	2005-06 NFHS			
		Urban		Rural	
		M	F	M	F
9.1.1	Abstinence of never-married young men and women. Ages 15-24.	90	99.6	86.8	99
9.1.1a	<i>Abstinence of never-married young men and women. Ages 15-19.</i>	M: 91.4		F: 99.4	
9.1.1b	<i>Abstinence of never-married young men and women. Ages 20-24.</i>	M: 83		F: 98.7	
9.1.2	Sex before the age of 15 Ages 15-24	1	4.5	3	12.7
9.1.3	Sex before the age of 18 Ages 18-24	6.4	24.3	14.8	48.1
9.2	Young people having premarital sex in last year. Ages 15-24.	5.6	0.3	7.3	0.7
9.2a	<i>Young people having premarital sex in last year. Ages 15-19.</i>	M: 5.1		F: 0.4	
9.2b	<i>Young people having premarital sex in last year. Ages 20-24.</i>	M: 8.9		F: 0.8	
9.3	Young people using a condom during premarital sex. Ages 15-24.	52.9	36.1	27.7	12.9
9.4	Young people having multiple partners last year. Ages 15-24.	M: 7		F: 0.1	
9.5	Young people using a condom at last higher risk sex. Ages 15-24.	M: 36.5		F: 22.2	
9.5.1	Young people using a condom at last higher risk sex of all young people surveyed.	53.1	36.5	28.2	17.4
9.6	Condom use at first sex. Ages 15-24.	25.5	4.1	11.3	2.3
9.7	Age-mixing in sexual relationships (young women age 15-19, non-marital, non-cohabiting partner in the last 12 months)	F: 11			

HIV Prevalence

#	Indicator	2005-06 NFHS			
		Urban		Rural	
		M	F	M	F
15.1	HIV prevalence among young people aged 15-24.	0.11	0.16	0.08	0.09
15.1a	<i>HIV prevalence among young people. Ages 15-19.</i>	0	0.08	0.01	0.06
15.1b	<i>HIV prevalence among young people. Ages 20-24.</i>	0.23	0.24	0.16	0.13
15.2	HIV prevalence among general population. Ages 15-49.	0.41	0.29	0.32	0.18
	HIV prevalence among general population. Ages 15-49. (sexes combined)	0.28			

Notes: Numbers indicate percentages unless otherwise stated
 Figures marked with an asterisk are based on small denominators (typically 25-49 unweighted cases).
 Figures with denominators less than 25 cases are not displayed.
 Bolded indicators are PEPFAR-specific indicators